



(levonorgestrel and ethinyl estradiol tablets, USP,
and ferrous bisglycinate tablets)
0.1mg/0.02mg and 36.5mg

BALCOLTRA SAVINGS PROGRAM COUPON

PAY AS LITTLE AS*:
\$10 PER 30-DAY FILL
\$15 PER 60-DAY FILL
\$20 PER 90-DAY FILL

BIN# 601341

RxGRP# OH5510041

PCN# OHCP

RxID# EE1100136187

Be sure to present this coupon, your insurance card (if applicable), and a valid prescription for Balcoltra to your participating pharmacist.

*For eligible patients. See full program terms and conditions below.

PHARMACIST INSTRUCTIONS

This coupon must be accompanied by a valid prescription for Balcoltra. Please submit the co-pay authorized by the patient's primary insurance as a secondary transaction to OPUS Health. For self-pay patients, submit the claim to U&C. **Pharmacists with questions, please call OPUS Health at 1-800-364-4767.**

Terms, Conditions, and Eligibility Criteria

This offer is only good for self-pay patients and commercial-prescription-drug-insured patients and is good for use only with a Balcoltra® (levonorgestrel and ethinyl estradiol tablets and ferrous bisglycinate tablets) prescription at the time the prescription is filled by the pharmacist and dispensed to the patient.

Depending on your insurance coverage, most eligible insured patients pay \$10 for a 30-day fill, \$15 for a 60-day fill, or \$20 for a 90-day fill. Patients may have additional financial responsibilities depending on your insurance plan.

This offer is not valid for patients enrolled in Medicare, Medicaid, TRICARE, or other federal or state programs (including any state pharmaceutical assistance programs), or private indemnity or HMO insurance plans that reimburse you for the entire cost of your prescription drugs. Patients may not use this coupon if they are Medicare-eligible and enrolled in an employer-sponsored health plan or prescription drug benefit program for retirees. Qualified health plans purchased through a health insurance exchange or marketplace established by a state government or the federal government are not considered to be government programs and are eligible for this program.

Offer not valid in any states where prohibited by law, taxed, or restricted.

The selling, purchasing, trading, or counterfeiting of this coupon is prohibited by law.

By redeeming this coupon, you acknowledge that you understand and agree to comply with the terms and conditions of this offer.

This coupon is the property of Avion Pharmaceuticals, LLC, and OPUS Health and must be returned upon request.

This coupon is not health insurance.

Each coupon is valid for unlimited refills. This coupon is not transferable.

Avion Pharmaceuticals, LLC reserves the right to rescind, revoke, or amend this offer at any time without notice.

Please see Important Risk Information on following page and full Prescribing Information, including Boxed Warning at balcoltra.com/pi

Questions about the Balcoltra Savings Program? Call 1-877-838-3846

INDICATION

What is Balcoltra?

Balcoltra is a prescription birth control pill used for the prevention of pregnancy.

IMPORTANT RISK INFORMATION, INCLUDING BOXED WARNING

WARNING TO WOMEN WHO SMOKE

Do not use Balcoltra if you smoke cigarettes and are over 35 years old. Smoking increases your risk of serious cardiovascular side effects (heart and blood vessel problems) from birth control pills, including death from heart attack, blood clots, or stroke. This risk increases with age and the number of cigarettes you smoke.

Who should not take Balcoltra?

Do not use Balcoltra if you have or have had blood clots, history of heart attack or stroke, high blood pressure that medicine cannot control, breast cancer or any cancer that is sensitive to female hormones, liver disease or liver tumors, unexplained bleeding from the vagina, hypersensitivity to any of the components, if you are or may be pregnant, or if you take Hepatitis C drugs containing ombitasvir/paritaprevir/ritonavir, with or without dasabuvir, as this may increase levels of liver enzymes in the blood.

What else should I know about taking Balcoltra?

Treatment with Balcoltra should be stopped if you have a blood clot, and at least 4 weeks before and through 2 weeks after major surgery. You should not take Balcoltra any earlier than 4 weeks after having a baby. If you are breastfeeding, talk to your doctor before taking Balcoltra. If you experience yellowing of the skin or eyes due to problems with your liver, you should stop taking Balcoltra. If you are prediabetic or diabetic, your doctor should monitor you while using Balcoltra. Your doctor should evaluate you if you have any significant change in headaches or irregular menstrual bleeding. Balcoltra contains FD&C Yellow No. 5 and may cause an allergic reaction, including in those with an allergy to aspirin.

What are the most serious risks of taking Balcoltra?

Balcoltra increases the risk of serious conditions, including blood clots, stroke, and heart attack. These can be life-threatening and require immediate medical care.

What are the possible side effects of Balcoltra?

The most common side effects of Balcoltra are headache, spotting or bleeding between periods or no menstrual bleeding, nausea, breast tenderness or pain, stomach pain, pain during periods, depression, acne, and vaginal infections.

Birth control pills do not protect you against any sexually transmitted disease, including HIV, the virus that causes AIDS.

You may report side effects to Avion Pharmaceuticals, LLC at 1-888-612-8466 or to the FDA at 1-800-FDA-1088.

Please see full Prescribing Information, including BOXED WARNING, at [balcoltra.com/pi](https://www.balcoltra.com/pi) or scan code

